

Package ‘MeTo’

January 11, 2019

Type Package

Title Meteorological Tools

Version 0.1.0

Maintainer Ullrich Dettmann <ullrich.dettmann@thuenen.de>

Description Meteorological Tools following the FAO56 irrigation paper of Allen et al. (1998) [1].

Functions for calculating:
reference evapotranspiration (ETref),
extraterrestrial radiation (Ra),
net radiation (Rn),
saturation vapor pressure (satVP),
global radiation (Rs),
soil heat flux (G),
daylight hours,
and more.

[1] Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56. FAO, Rome, 300(9).

BugReports <https://bitbucket.org/Ullid/meto/issues>

Imports lubridate

License GPL (>= 2)

Depends R (>= 3.3.0)

Encoding UTF-8

LazyData true

RoxygenNote 6.1.1

Suggests testthat

NeedsCompilation no

Author Ullrich Dettmann [aut, cre],
R. Grimma [aut]

Repository CRAN

Date/Publication 2019-01-11 17:00:16 UTC

R topics documented:

| | |
|---------------------------|-----------|
| adj_u2 | 2 |
| constDefaults | 3 |
| controlDefaults | 4 |
| deltaVP | 5 |
| dlh | 6 |
| dr | 7 |
| e0 | 7 |
| estG | 8 |
| estP | 10 |
| estRs | 11 |
| ETref | 12 |
| prep.date | 15 |
| psyc_cons | 15 |
| Ra | 16 |
| Rn | 18 |
| Rnl | 20 |
| Rns | 22 |
| Rso | 23 |
| satVP | 24 |
| SolarDec | 25 |
| VP | 26 |
| w | 27 |
| ws | 28 |
| Index | 29 |

| | |
|--------|-------------------------------|
| adj_u2 | <i>Adjust wind speed data</i> |
|--------|-------------------------------|

Description

Adjust wind speed data to 2 meter height.

Usage

```
adj_u2(u, uz)
```

Arguments

| | |
|----|--|
| u | measured wind speed at uz above ground surface [m/s] |
| uz | height of windspeed measurement above ground surface [m] |

Note

eq. 47 of reference

References

Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56. FAO, Rome, 300(9).

Examples

```
adj_u2(3.2, uz = 10)
```

constDefaults

Constants

Description

Constants required for calculations.

Usage

```
constDefaults
```

Format

GSC: solar constant (0.0820 [MJ/(m² min)])

sigma: Stefan-Boltzmann constant (4.903 10⁻⁹ [MJ/(K⁴ m² day)])

a1: constant lapse rate moist air (0.0065 [K/m]) (e.g. in [estP](#))

R: specific gas constant (287 [J/(kg K)]) (e.g. in [estP](#))

g: gravitational acceleration (9.807 [m/s²]) (e.g. in [estP](#))

eps: ratio molecular weight of water vapour/dry air (0.622) (e.g. in [psyc_cons](#))

lambda: latent heat of vaporization (2.45 [MJ/kg]) (e.g. in [psyc_cons](#))

cp: specific heat at constant pressure (1.013 10⁻³ [MJ/(kg degreeC)]) (e.g. in [psyc_cons](#))

| | |
|-----------------|-------------------------------|
| controlDefaults | <i>Control default values</i> |
|-----------------|-------------------------------|

Description

Default values for control. If necessary modify with `control = list()` in function call.

Usage

`controlDefaults`

Format

albedo: 0.23 [-] for hypothetical grass and alfalfa reference crops used in the FAO-56 PM equations (e.g. in [ETref](#), [Rn](#) or [estG](#))

Po: atmospheric pressure at sea level (101.3 [kPa])

z0: elevation at reference level (0 [m]) (e.g. in [ETref](#) or [estP](#))

Tko: reference temperature [degreeC] at elevation z0. Often assumed to be 20 degreeC (e.g. in [ETref](#) or [estP](#))

uz: height of windspeed measurement above ground surface (2 [m]) (e.g. in [ETref](#) or [adj_u2](#))

Lz:

longitude of the centre of the local time zone (degrees west of Greenwich)

- 0 for Greenwich

- 345 for Germany

- 330 for Cairo (Egypt)

- 255 for Bangkok (Thailand)

- 75, 90, 105 and 120 for Eastern, Central, Rocky Mountain and Pacific time zones (United States)

Lz is only needed if calculation period is shorter 1 day.

as: regression constant, expressing fraction of extraterrestrial radiation reaching earth on overcast days (default = 0.25) (e.g. in [ETref](#), [Rn](#) or [estRs](#))

bs: $a_s + b_s$ fraction of extraterrestrial radiation reaching earth on clear days (default = 0.5) (e.g. in [ETref](#), [Rn](#) or [estRs](#))

est.ratio.Rs.Rso:

R_s/R_{s0} is used to represent cloud cover. For hourly or shorter periods during the nighttime, the ratio R_s/R_{s0} is set equal to the R_s/R_{s0} calculated for a time period occurring 2-3 hours before sunset. If single values during nighttime are calculated R_s/R_{s0} ratio 2-3 hours before sunset can not be calculated and an approximation is needed. Following Allen (1999) one can assume $R_s/R_{s0} = 0.4$ to 0.6 during nighttime periods in humid and subhumid climates and $R_s/R_{s0} = 0.7$ to 0.8 in arid and semiarid climates. A value of $R_s/R_{s0} = 0.3$ presumes total cloud cover.

References

Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56. FAO, Rome, 300(9).

 deltaVP

Slope of the saturation vapor pressure curve

Description

Slope of the saturation vapor pressure curve [kPa/degreeC].

Usage

deltaVP(Tmean = NULL, Tmax = NULL, Tmin = NULL)

Arguments

| | |
|-------|--|
| Tmean | Mean Temperature [degreeC] (mean daily, mean hourly, etc. air temperature) |
| Tmax | maximum temperature during 24-hour period [degreeC] (if Tmean is missing) |
| Tmin | minimum temperature during 24-hour period [degreeC] (if Tmean is missing) |

Details

valid for daily, hourly and shorter periods

Value

slope of the saturation vapor pressure curve [kPa/degreeC]

Note

eq. 13 of reference

References

Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56. FAO, Rome, 300(9).

Examples

```
deltaVP(Tmax = 34.8, Tmin = 25.6)
deltaVP(Tmean = 30.2)
```

| | |
|-----|----------------------------|
| dlh | <i>Daylighth hours (N)</i> |
|-----|----------------------------|

Description

Daylighth hours in dependence to latitude.

Usage

```
dlh(x, lat.rad = NULL, lat.deg = NULL)
```

Arguments

| | |
|---------|--|
| x | date-time object or day of the year |
| lat.rad | latitude [rad] (either lat.rad or lat.deg). Latitude is positive for the northern hemisphere and negative for the southern hemisphere |
| lat.deg | latitude [degree] (either lat.rad or lat.deg). Latitude is positive for the northern hemisphere and negative for the southern hemisphere |

Details

x: must be provided as number (1-366) or as a common date-time object (e.g, POSIXct, POSIXlt or Date objects). All formats for which is.timepoint from the lubridate package returns TRUE can be used

Note

eq. 34 of reference

References

Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56. FAO, Rome, 300(9).

Examples

```
dlh(x = 105, lat.deg = 13.73)
dlh(x = 105, lat.rad = 0.283)
dlh(x = as.Date('2018-04-15'), lat.deg = 13.73)
```

dr *Inverse relative distance Earth-Sun*

Description

Calculate inverse relative distance Earth-Sun for daily and shorter periods.

Usage

dr(x)

Arguments

x date-time object or Day of the year

Details

x: must be provided as number (1-366) or as a common date-time object (e.g, POSIXct, POSIXlt or Date objects). All formats for which is.timepoint from the lubridate package returns TRUE can be used

Note

eq. 23 of reference

References

Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56. FAO, Rome, 300(9).

e0 *Saturation Vapor Pressure*

Description

Mean saturation vapour pressure [kPa] for short time intervals less than a day. Calculated with air temperature.

Usage

e0(Temp)

Arguments

Temp Temperature [degreeC]

Value

saturation vapour pressure at air temperature [kPa/degreeC])

Note

eq. 11 of the reference

for day, week, decade or month, the mean saturation vapour pressure should be computed with [satVP](#)

References

Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56. FAO, Rome, 300(9).

See Also

[satVP](#), [VP](#)

estG

Estimate soil heat flux (G)

Description

Estimate soil heat flux (G) for periods shorter than a day.

Usage

```
estG(x, Rs, Tmean, Rhmean, lat.rad = NULL, lat.deg = NULL,
     long.deg = NULL, elev = 1, t1, control = list(albedo = 0.23, Lz =
     345, est.ratio.Rs.Rso = NA))
```

Arguments

| | |
|----------|---|
| x | date-time object (see details) |
| Rs | incoming solar radiation [MJ/(m2 time)] |
| Tmean | Mean air temperature [degreeC] |
| Rhmean | Mean air humidity [percent] |
| lat.rad | latitude [rad]. Use either lat.rad or lat.deg. Latitude is positive for the northern hemisphere and negative for the southern hemisphere |
| lat.deg | latitude [degree]. Use either lat.deg or lat.rad. Latitude is positive for the northern hemisphere and negative for the southern hemisphere |
| long.deg | longitude of the measurement site (degrees east of Greenwich) |
| elev | station elevation above sea level [m] |

| | |
|---------|--|
| t1 | length of calculation period [hour] (1 for hourly period, 0.5 for a 30-minute period or 24 for daily period). Only needed if x is date-time object with length of 1. |
| control | list for control parameters and empirical factors defined in <code>controlDefaults</code> and <code>constDefaults</code> (see Details) |

Details

during daylight periods G is estimated to be $R_n \times 0.1$ During nighttime $G = R_n \times 0.5$. Day is defined for extraterrestrial radiation > 0 .

x: must be provided as a common date-time object (e.g, POSIXct, POSIXlt or Date objects). All formats for which `is.timepoint` from the `lubridate` package returns TRUE can be used

control:

albedo: default 0.23 for the hypothetical grass and alfalfa reference crops used in the FAO-56 PM equations

Lz:

longitude of the centre of the local time zone (degrees west of Greenwich)

- 0 for Greenwich

- 345 for Germany

- 330 for Cairo (Egypt)

- 255 for Bangkok (Thailand)

- 75, 90, 105 and 120 for Eastern, Central, Rocky Mountain and Pacific time zones (United States)

Lz is only needed if calculation period is shorter 1 day.

est.ratio.Rs.Rso: Rs/Rso is used to represent cloud cover. For hourly periods during the nighttime, the ratio Rs/Rso is set equal to the Rs/Rso calculated for a time period occurring 2-3 hours before sunset. If single values during nighttime are calculated Rs/Rso ration 2-3 hours before sunset can not be calculated and an approximation is needed. Following Allen (1999) one can assume $R_s/R_{s0} = 0.4$ to 0.6 during nighttime periods in humid and subhumid climates and $R_s/R_{s0} = 0.7$ to 0.8 in arid and semiarid climates. A value of $R_s/R_{s0} = 0.3$ presumes total cloud cover.

Note

eq. 45 and 46 of reference

References

Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56. FAO, Rome, 300(9).

Examples

```
estG(x = as.POSIXct(c('2018-10-01 14:30', '2018-10-01 15:00')), Tmean = 38, Rhmean = 52, Rs = 2.450,
     elev = 8, lat.deg = 16.21, long.deg = 343.75, control = list(Lz = 15))
```

```
estG(x = as.POSIXct('2018-10-01 02:30'), Tmean = 28, Rhmean = 90, t1 = 1, Rs = 0, elev = 8,
```

```
lat.deg = 16.2, long.deg = 343.75, control = list(Lz = 15, est.ratio.Rs.Rso = 0.8))
estG(x = as.POSIXct('2018-10-01 14:30'), Tmean = 38, Rhmean = 52, t1 = 1, Rs = 2.450, elev = 8,
lat.deg = 16.21, long.deg = 343.75, control = list(Lz = 15))
```

estP

Estimate atmospheric pressure (P)

Description

Values for atmospheric pressure as a function of altitude.

Usage

```
estP(elev, control = list(Tko = 20))
```

Arguments

| | |
|---------|--|
| elev | elevation [m] |
| control | list for control parameters and empirical factors defined in controlDefaults and constDefaults (see Details) |

Details

Control variables:

Tko: reference temperature [degreeC] at elevation z0. Often assumed to be 20 degreeC

z0: elevation at reference level [m]

a1: constant lapse rate moist air (0.0065 [K/m])

g: gravitational acceleration (9.807 [m/s²])

R: specific gas constant (287 [J/(kg K)])

Value

atmospheric pressure [kPa]

Note

eq. 3-2 of Reference

References

Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56. FAO, Rome, 300(9).

Examples

```
estP(elev = 25, control = list(Tko = 20))
```

estRs *Estimate solar radiation (Rs)*

Description

Rs is calculated from relative sunshine duration and extraterrestrial radiation with the Angstrom formula.

Usage

```
estRs(x, n, lat.rad = NULL, lat.deg = NULL, t1, control = list(as =
  0.25, bs = 0.5))
```

Arguments

| | |
|---------|--|
| x | date-time object or Day of the year |
| n | actual duration of sunshine [hour] |
| lat.rad | latitude [rad] (either lat.rad or lat.deg). Latitude is positive for the northern hemisphere and negative for the southern hemisphere |
| lat.deg | latitude [degree] (either lat.rad or lat.deg). Latitude is positive for the northern hemisphere and negative for the southern hemisphere |
| t1 | length of calculation period [hour] (1 for hourly, 0.5 for a 30-minute or 24 for daily period). Only needed if length of x is date-time object with length of 1. |
| control | list for control parameters and empirical factors defined in controlDefaults and constDefaults (see Details) |

Details

control:

as: regression constant, expressing fraction of extraterrestrial radiation reaching earth on overcast days ($n = 0$) (default = 0.25)

bs: as + bs fraction of extraterrestrial radiation reaching earth on clear days ($n = N$) (default = 0.5)

x: must be provided as.numeric (1-366) or as a common date-time object (e.g, POSIXct, POSIXlt or Date objects). All formats for which is.timepoint from the lubridate package returns TRUE can be used

Value

solar or shortwave radiation (Rs) [MJ/(m² day)]

Note

eq. 35 of reference

References

Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56. FAO, Rome, 300(9).

Examples

```
estRs(x = 135, n = 7, lat.rad = NULL, lat.deg = -22.9)
```

ETref

FAO-56 Penman-Monteith grass reference evapotranspiration

Description

FAO Penman-Monteith equations to compute grass reference evapotranspiration from weather data for daily, monthly, hourly or shorter periods.

Usage

```
ETref(x, Tmax = NULL, Tmin = NULL, Rhmax = NULL, Rhmin = NULL,
      Tmean = NULL, Rhmean = NULL, u = NULL, Rs = NULL, n = NULL,
      P = NULL, elev, lat.rad = NULL, lat.deg = NULL, long.deg = NULL,
      t1, G = NULL, actVP = NULL, control = list())
```

Arguments

| | |
|---------|--|
| x | date-time object or day of the year (must be date-time object if calculation period is shorter than a day) |
| Tmax | maximum temperature [degreeC] during 24-hour period (for daily values) |
| Tmin | minimum temperature [degreeC] during 24-hour period (for daily values) |
| Rhmax | maximum of air humidity [percent] during 24-hour period (for daily values) |
| Rhmin | minimum of air humidity [percent] during 24-hour period(for daily values) |
| Tmean | mean air temperature [degreeC]. For periods shorter 1 day. |
| Rhmean | mean relative air humidity [percent]. For periods shorter 1 day or if Rhmax and Rhmin are missing. |
| u | wind speed [m/s] at 2 meter height. If measurement height is not 2 m, define height with control <- list(uz = 2) |
| Rs | solar radiation [MJ/(m2 time)] |
| n | actual hours of sunshine. Used to calculate Rs if Rs measurements are not available (see estRs). |
| P | air pressure [kPa]. Estimated with estP if missing. |
| elev | station elevation above sea level [m] |
| lat.rad | latitude [rad]. Use either lat.rad or lat.deg. Latitude is positive for the northern hemisphere and negative for the southern hemisphere |

| | |
|----------|---|
| lat.deg | latitude [degree]. Use either lat.deg or lat.rad. Latitude is positive for the northern hemisphere and negative for the southern hemisphere |
| long.deg | longitude [degree] east of Greenwich (for periods < 1 day) |
| t1 | length of calculation period [hour] (1 for hourly, 0.5 for 30-minute and 24 for daily period). Only needed if x is date-time object with length of 1. |
| G | soil heat flux (Assumed to be 0 for daily calculations) (for calculation periods shorter than a day estimated with estG if missing) |
| actVP | Actual vapor pressure [kPa]. If Rhmax, Rhmin and Rhmean are NULL |
| control | list for control parameters and empirical factors (see details, controlDefaults and constDefaults) |

Details

x: must be provided as.numeric (1-366) or as a common date-time object (e.g, POSIXct, POSIXlt or Date objects). All formats for which is.timepoint from the lubridate package returns TRUE can be used

control: (see also [controlDefaults](#) and [constDefaults](#)) Lz:

longitude of the centre of the local time zone (degrees west of Greenwich)

- 0 for Greenwich

- 345 for Germany

- 330 for Cairo (Egypt)

- 255 for Bangkok (Thailand)

- 75, 90, 105 and 120 for Eastern, Central, Rocky Mountain and Pacific time zones (United States)

Lz is only needed if calculation period is shorter than 1 day.

uz: height of wind measurements (m)

albedo: default 0.23 for the hypothetical grass and alfalfa reference crops used in the FAO-56 PM equations

as: regression constant, expressing fraction of extraterrestrial radiation reaching earth on overcast days (n = 0) (default = 0.25)

bs: as + bs fraction of extraterrestrial radiation reaching earth on clear days (n = N) (default = 0.5)

Tko: reference temperature [degreeC] at elevation z0. Only needed if atmospheric pressure is missing. Often assumed to be 20 degreeC.

z0: elevation at reference level (default = 0 [m])

est.ratio.Rs.Rso:

Rs/Rso is used to represent cloud cover. For hourly or shorter periods during the nighttime, the ratio Rs/Rso is set equal to the Rs/Rso calculated for a time period occurring 2-3 hours before sunset. If single values during nighttime are calculated Rs/Rso ration 2-3 hours before

sunset can not be calculated and an approximation is needed. Following Allen (1999) one can assume $R_s/R_{so} = 0.4$ to 0.6 during nighttime periods in humid and subhumid climates and $R_s/R_{so} = 0.7$ to 0.8 in arid and semiarid climates. A value of $R_s/R_{so} = 0.3$ presumes total cloud cover.

Value

grass reference evapotranspiration [mm]

Note

eq. 6 from reference for daily and eq. 53 for hourly or shorter periods

Author(s)

Ulrich Dettmann

References

Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56. FAO, Rome, 300(9).

Examples

```
# -----
# Daily Evapotranspiration
# -----

ETref(x = 187, Rs = 22.07, elev = 100, lat.deg = 50.8, Tmax = 21.5, Tmin = 12.3,
      Rhmax = 84, Rhmin = 63,
      u = 2.78, control = list(uz = 10), P = 100.1)

# Calculation with sunshine hour (n) instead of
# global radiation (Rs) (Rs ist estimated from n with estRs):

ETref(x = 187, n = 9.25, elev = 100, lat.deg = 50.8, Tmax = 21.5, Tmin = 12.3,
      Rhmax = 84, Rhmin = 63,
      u = 2.78, control = list(uz = 10), P = 100.1)

# -----
# Hourly Evapotranspiration
# -----

ETref(x = as.POSIXct(c('2018-10-01 14:30', '2018-10-01 15:30')),
      Tmean = c(38, 37.8), Rhmean = c(52, 52.3), u = c(3.3, 3.2), Rs = c(2.450, 2.5), elev = 8,
      lat.deg = 16.22, long.deg = 343.75, G = c(0.175, 0.178) , P = c(101.21, 101.21) ,
      control = list(Lz = 15))

# If only one time step is calculated t1 must be provided (1 for hourly, 0.5 for 30 minute periods):

ETref(x = as.POSIXct('2018-10-01 14:30'), t1 = 1,
      Tmean = 38, Rhmean = 52, u = 3.3, Rs = 2.450, elev = 8,
```

```

lat.deg = 16.22, long.deg = 343.75, G = 0.1749218, P = 101.2056,
control = list(Lz = 15))

# Calculation with missing soil heat flux (G) and atmospheric pressure (P) (G is estimated with estG
# and P with estP)

ETref(x = as.POSIXct('2018-10-01 14:30'), t1 = 1,
      Tmean = 38, Rhmean = 52, u = 3.3, Rs = 2.450, elev = 8,
      lat.deg = 16.22, long.deg = 343.75,
      control = list(Lz = 15))
# -----

```

| | |
|-----------|---------------------|
| prep.date | <i>Prepare date</i> |
|-----------|---------------------|

Description

Checks if x is date-time object or day of the year (doy) and returns doy.

Usage

```
prep.date(x)
```

Arguments

| | |
|---|-------------------------------------|
| x | date-time object or Day of the year |
|---|-------------------------------------|

Value

Day of the year

| | |
|-----------|-------------------------------|
| psyc_cons | <i>psychrometric constant</i> |
|-----------|-------------------------------|

Description

The psychrometric constant [kPa/degreeC]

Usage

```
psyc_cons(elev, P = NULL, control = list())
```

Arguments

| | |
|---------|--|
| elev | elevation [m] |
| P | atmospheric pressure [kPa] |
| control | list for control parameters and empirical factors defined in controlDefaults and constDefaults (see Details) |

Details

control: Tko: reference temperature [degreeC] at elevation z0. Only needed if atmospheric pressure is missing. Often assumed to be 20 degreeC

elev: station elevation above sea level [m]. Needed if P = NULL for `estP`

lambda: latent heat of vaporization [MJ/kg]

eps: ratio molecular weight of water vapor/dry air = 0.622

cp: specific heat of moist air = 1.013×10^{-3} [MJ/(kg degreeC)]

Value

psychrometric constant [kPa/degreeC]

Note

eq. 8 of reference

References

Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56. FAO, Rome, 300(9).

Examples

```
psyc_cons(elev = 2, P = 101.3)
```

 Ra

Extraterrestrial radiation

Description

Extraterrestrial radiation [MJ/(m2 time)] in dependence to time, latitude and longitude.

Usage

```
Ra(x, lat.rad = NULL, lat.deg = NULL, long.deg, t1, control = list(Lz
= 345))
```

Arguments

| | |
|----------|--|
| x | date-time object or day of the year (must be date-time object if calculation period is smaller 1 day) |
| lat.rad | latitude [rad]. Use either lat.rad or lat.deg. Latitude is positive for the northern hemisphere and negative for the southern hemisphere |
| lat.deg | latitude [degree]. Use either lat.deg or lat.rad. Latitude is positive for the northern hemisphere and negative for the southern hemisphere |
| long.deg | longitude of the measurement site (degrees east of Greenwich) (only needed for periods shorter 1 day) |
| tl | length of calculation period [hour] (1 for hourly period, 0.5 for a 30-minute period or 24 for daily period). Only needed if x is date-time object with length of 1. |
| control | list for control parameters and empirical factors defined in controlDefaults and constDefaults (see Details) |

Details

x: must be provided as.numeric (1-366) or as a common date-time object (e.g, POSIXct, POSIXlt or Date objects). All formats for which is.timepoint from the lubridate package returns TRUE can be used

control:

Lz (for periods < 1 day):

longitude of the centre of the local time zone (degrees west of Greenwich)

- 0 for Greenwich

- 345 for Germany

- 330 for Cairo (Egypt)

- 255 for Bangkok (Thailand)

- 75, 90, 105 and 120 for Eastern, Central, Rocky Mountain and Pacific time zones (United States)

Note

eq. 21 (period = 1 day) and eq. 28 (hourly or shorter) of the reference

References

Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56. FAO, Rome, 300(9).

Examples

```
# -----
# Daily period
# -----

Ra(x = 105, lat.deg = 13.73)
```

```
# -----
# Hourly period
# -----

Ra(x = as.POSIXct(c('2018-10-01 14:30', '2018-10-01 15:30')),
   lat.deg = 16.21, long.deg = 343.75, control = list(Lz = 15))

Ra(x = as.POSIXct('2018-10-01 14:30'), t1 = 1,
   lat.deg = 16.21, long.deg = 343.75, control = list(Lz = 15))
```

Rn *Net radiation (Rn)*

Description

Difference between the incoming net shortwave radiation ([Rns](#)) and the outgoing net longwave radiation ([Rnl](#)).

Usage

```
Rn(x, Tmax = NULL, Tmin = NULL, Rhmax = NULL, Rhmin = NULL,
   Rs = NULL, n = NULL, elev, lat.rad = NULL, lat.deg = NULL,
   long.deg = NULL, Rhmean = NULL, actVP = NULL, Tmean = NULL, t1,
   control = list())
```

Arguments

| | |
|----------|---|
| x | date-time object or day of the year (must be date-time object if calculation period is shorter than a day) |
| Tmax | maximum temperature [degreeC] during 24-hour period (for daily values) |
| Tmin | minimum temperature [degreeC] during 24-hour period (for daily values) |
| Rhmax | daily maximum of air humidity [percent] (for daily values) |
| Rhmin | daily minimum of air humidity [percent] (for daily values) |
| Rs | incoming solar radiation [MJ/(m2 time)] |
| n | Actual hours of sunshine. Used to calculate Rs if missing. |
| elev | station elevation above sea level [m] |
| lat.rad | latitude [rad]. Use either lat.rad or lat.deg. Latitude is positive for the northern hemisphere and negative for the southern hemisphere |
| lat.deg | latitude [degree]. Use either lat.deg or lat.rad. Latitude is positive for the northern hemisphere and negative for the southern hemisphere |
| long.deg | longitude of the measurement site (degrees east of Greenwich) (for periods < 1 day) |
| Rhmean | Mean air humidity [percent] for periods < day or if Rhmax and Rhmin are missing |

| | |
|---------|--|
| actVP | Actual vapor pressure [kPa]. If Rhmax, Rhmin and Rhmean are NULL |
| Tmean | Mean air temperature [degree C] for periods < day |
| t1 | length of calculation period [hour] (1 for hourly period, 0.5 for a 30-minute period or 24 for daily period). Only needed if x is date-time object with length of 1. |
| control | list for control parameters and empirical factors (see Details) |

Details

for daily and hourly calculations

x: must be provided as numeric (1-366) or as a common date-time object (e.g, POSIXct, POSIXlt or Date objects). All formats for which is.timepoint from the lubridate package returns TRUE can be used

control:

albedo: default 0.23 for the hypothetical grass and alfalfa reference crops used in the FAO-56 PM equations

as: regression constant, expressing fraction of extraterrestrial radiation reaching earth on overcast days (n = 0) (default = 0.25)

bs: as + bs fraction of extraterrestrial radiation reaching earth on clear days (n = N) (default = 0.5)

Value

net radiation

Note

eq. 40 of reference

References

Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56. FAO, Rome, 300(9).

See Also

[Rns](#), [Rn1](#)

Examples

```
# -----
# Daily period
# -----

Rn(x = 105, n = 8.5, elev = 2, actVP = 2.85, Tmax = 34.8,
   Tmin = 25.6, lat.deg = 13.73)

Rn(x = 135, elev = 1, Rs = 14.5, Tmax = 25.1, Tmin = 19.1,
```

```

    lat.deg = -22.9, actVP = 2.1)

# -----
# Hourly period
# -----

Rn(x = as.POSIXct(c('2018-10-01 14:30', '2018-10-01 15:30')), Tmean = c(38, 37.8),
   Rhmean = c(52, 52.2), Rs = c(2.450, 2.1), elev = 8, lat.deg = 16.2,
   long.deg = 343.75, control = list(Lz = 15))

Rn(x = as.POSIXct('2018-10-01 14:30'), Tmean = 38, Rhmean = 52, t1 = 1, Rs = 2.450,
   elev = 8, lat.deg = 16.2, long.deg = 343.75, control = list(Lz = 15))

```

Rnl

Net longwave radiation (Rnl)

Description

Net longwave radiation (Rnl).

Usage

```

Rnl(x, Tmax = NULL, Tmin = NULL, Rhmax = NULL, Rhmin = NULL,
    Rs = NULL, lat.rad = NULL, lat.deg = NULL, long.deg = NULL, elev,
    actVP = NULL, t1, Tmean = NULL, Rhmean = NULL, control = list())

```

Arguments

| | |
|----------|---|
| x | date-time object or day of the year (must be date-time object if calculation period is shorter than a day) |
| Tmax | maximum temperature [degreeC] during 24-hour period (for daily values) |
| Tmin | minimum temperature [degreeC] during 24-hour period (for daily values) |
| Rhmax | daily maximum of air humidity [percent] (for daily values) |
| Rhmin | daily minimum of air humidity [percent] (for daily values) |
| Rs | incoming solar radiation [MJ/(m2 time)] |
| lat.rad | latitude [rad]. Use either lat.rad or lat.deg. Latitude is positive for the northern hemisphere and negative for the southern hemisphere |
| lat.deg | latitude [degree]. Use either lat.deg or lat.rad. Latitude is positive for the northern hemisphere and negative for the southern hemisphere |
| long.deg | see Rso |
| elev | station elevation above sea level [m] |
| actVP | Actual vapor pressure [kPa]. If Rhmax, Rhmin and Rhmean are NULL |

| | |
|---------|--|
| t1 | length of calculation period [hour] (1 for hourly period, 0.5 for a 30-minute period or 24 for daily period). Only needed if length of x is date-time object with length of 1. |
| Tmean | mean temperature [degreeC] during the time period (for periods shorter than a day) |
| Rhmean | Mean air humidity [percent] (for periods shorter than a day or if Rhmax and Rhmin are missing) |
| control | list for control parameters and empirical factors (see Details and controlDefaults) |

Details

x: must be provided as.numeric (1-366) or as a common date-time object (e.g, POSIXct, POSIXlt or Date objects). All formats for which is.timepoint from the lubridate package returns TRUE can be used

control: Lz:

longitude of the centre of the local time zone (degrees west of Greenwich)

- 0 for Greenwich

- 345 for Germany

- 330 for Cairo (Egypt)

- 255 for Bangkok (Thailand)

- 75, 90, 105 and 120 for Eastern, Central, Rocky Mountain and Pacific time zones (United States)

Lz is only needed if calculation period is shorter 1 day.

est.ratio.Rs.Rso:

Rs/Rso is used to represent cloud cover. For hourly or shorter periods during the nighttime, the ratio Rs/Rso is set equal to the Rs/Rso calculated for a time period occurring 2-3 hours before sunset. If single values during nighttime are calculated Rs/Rso ration 2-3 hours before sunset can not be calculated and an approximation is needed. Following Allen (1999) one can assume Rs/Rso = 0.4 to 0.6 during nighttime periods in humid and subhumid climates and Rs/Rso = 0.7 to 0.8 in arid and semiarid climates. A value of Rs/Rso = 0.3 presumes total cloud cover.

Value

Rnl net outgoing longwave radiation [MJ/(m2 time)]

Note

eq. 39 of reference

References

Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56. FAO, Rome, 300(9).

Rns *Net solar or net shortwave radiation (Rns)*

Description

Net shortwave radiation is the balance between incoming and reflected solar radiation.

Usage

```
Rns(Rs, control = list(albedo = 0.23))
```

Arguments

| | |
|---------|---|
| Rs | incoming solar radiation [MJ/(m ² time)] |
| control | list for control parameters and empirical factors (see Details) |

Details

control:

albedo [-]: 0.23 for hypothetical grass and alfalfa reference crops used in the FAO-56 PM equations valid for daily and shorter periods

Value

Rns net solar or shortwave radiation [MJ/(m² time)]

Note

eq. 38 of reference

References

Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56. FAO, Rome, 300(9).

Examples

```
Rns(Rs = 22.1)
Rns(Rs = 22.1, control = list(albedo = 0.20))
```

Rso *Clear-sky solar radiation (Rso)*

Description

Clear-sky solar radiation for daily and shorter periods.

Usage

```
Rso(x, lat.rad = NULL, lat.deg = NULL, long.deg = NULL, elev, t1,
    control = list(Lz = 345))
```

Arguments

| | |
|----------|--|
| x | date-time object or day of the year (must be date-time object if calculation period is shorter than a day) |
| lat.rad | latitude [rad]. Use either lat.rad or lat.deg. Latitude is positive for the northern and negative for the southern hemisphere |
| lat.deg | latitude [degree]. Use either lat.deg or lat.rad. Latitude is positive for the northern and negative for the southern hemisphere |
| long.deg | longitude of the measurement site (degrees east of Greenwich) (only needed for periods < 1 day) |
| elev | station elevation above sea level [m] |
| t1 | length of calculation period [hour] (1 for hourly period, 0.5 for a 30-minute period or 24 for daily period). |
| control | list for control parameters and empirical factors defined in controlDefaults and constDefaults (see Details) |

Details

x: must be provided as.numeric (1-366) or as a common date-time object (e.g, POSIXct, POSIXlt, and Date objects). All formats for which is.timepoint from the lubridate package returns TRUE can be used

control:

Lz:

longitude of the centre of the local time zone (degrees west of Greenwich)

- 0 for Greenwich
- 345 for Germany
- 330 for Cairo (Egypt)
- 255 for Bangkok (Thailand)
- 75, 90, 105 and 120 for Eastern, Central, Rocky Mountain and Pacific time zones (United States)

Lz is only needed if calculation period is shorter 1 day.

for day, hour and shorter periods

References

eq. 37; Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56. FAO, Rome, 300(9).

Examples

```
# -----
# Daily period
# -----

Rso(x = 135, elev = 1, lat.deg = -22.9)

# -----
# Hourly period
# -----

Rso(x = as.POSIXct('2018-10-01 12:30'), t1 = 1, elev = 8, lat.deg = 16.2,
    long.deg = 343.75, control = list(Lz = 15))
```

satVP

Saturation Vapor Pressure

Description

Saturation vapor pressure [kPa].

Usage

```
satVP(Tmax = NULL, Tmin = NULL, Tmean = NULL, interval = "day",
      print.warning = T)
```

Arguments

| | |
|---------------|--|
| Tmax | maximum temperature [degreeC] for daily, weekly, monthly periods |
| Tmin | minimum temperature [degreeC] for daily, weekly, monthly periods |
| Tmean | mean air temperature [degreeC] for hourly or shorter periods |
| interval | hour, day, week or month |
| print.warning | TRUE or FALSE |

Details

interval: hour (eq. 11 of reference)

interval: day, week or month (eq. 12 of reference or eq. 11 if only Tmean is provided)

Value

Saturation Vapor Pressure [kPa]

References

Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56. FAO, Rome, 300(9).

See Also

[e0](#), [VP](#)

Examples

```
satVP(Tmax = 24.5, Tmin = 15, interval = 'day')
satVP(Tmax = 24.5, Tmin = 15, interval = 'week')
satVP(Tmax = 24.5, Tmin = 15, interval = 'month')

satVP(Tmax = 24.5, Tmin = 15, interval = 'hour')
satVP(Tmean = 19.75, interval = 'hour')
```

SolarDec

Solar declination

Description

Calculate solar declination for daily and shorter periods.

Usage

SolarDec(x)

Arguments

x date or day of the year

Details

x: must be provided as number (1-366) or as a common date-time object (e.g, POSIXct, POSIXlt or Date objects). All formats for which is.timepoint from the lubridate package returns TRUE can be used

Note

eq. 24 of reference

References

Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56. FAO, Rome, 300(9).

VP *Actual vapor pressure*

Description

Calculate actual vapour pressure [kPa] either from psychrometric data or from relative humidity data.

Usage

```
VP(Tmax = NULL, Tmin = NULL, Rhmax = NULL, Rhmin = NULL,
   interval = "day", Tmean = NULL, Rhmean = NULL, Twet = NULL,
   Tdry = NULL, apsy = NULL, P = NULL)
```

Arguments

| | |
|----------|--|
| Tmax | maximum temperature during 24-hour period (for daily values) |
| Tmin | minimum temperature during 24-hour period (for daily values) |
| Rhmax | maximum relative humidity [percent] (for daily values) |
| Rhmin | minimum relative humidity [percent] (for daily values) |
| interval | hour, day, week or month |
| Tmean | Mean air temperature [degreeC] (for periods shorter 1 day) |
| Rhmean | Mean air humidity [percent] (for periods shorter 1 day or if Rhmax and Rhmin are missing) |
| Twet | wet bulb temperature (for calculation with psychrometric data) |
| Tdry | dry bulb temperature (for calculation with psychrometric data) |
| apsy | coefficient depending on the type of ventilation of the wet bulb [kPa/(degreeC)] (for calculation with psychrometric data) |
| P | atmospheric pressure [kPa] |

Details

x: must be provided as.numeric (1-366) or as a common date-time object (e.g, POSIXct, POSIXlt or Date objects). All formats for which is.timepoint from the lubridate package returns TRUE can be used

interval: - use hour for periods <= one hour
 - for day, week or month the same equations are used

Note

eq. 17 of reference (Determination of actual vapour pressure from relative maximum and minimum humidity)

eq. 15 of reference (Actual vapour pressure derived from psychrometric data) (see [psyc_cons](#))

eq. 19 of reference (used in the absence of RHmax and RHmin)

eq. 54 of reference (for periods shorter than a day)

References

Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56. FAO, Rome, 300(9).

See Also

[e0](#), [satVP](#)

Examples

```
VP(Tmax = 25, Tmin = 18, Rhmax = 82, Rhmin = 54)
VP(Tmax = 25, Tmin = 18, Rhmean = 68)
```

w

Solar time angle

Description

Solar time angle at midpoint of the period (for periods < 1 day).

Usage

```
w(x, long.deg, control = list(Lz = 345))
```

Arguments

| | |
|----------|---|
| x | date-time object (e.g, POSIXct, POSIXlt or Date objects). |
| long.deg | longitude of the measurement site (degrees east of Greenwich) (for periods < 1 day) |
| control | list for control parameters and empirical factors (see Details) |

Details

control:

Lz:

longitude of the centre of the local time zone (degrees west of Greenwich)

- 0 for Greenwich
- 345 for Germany
- 330 for Cairo (Egypt)
- 255 for Bangkok (Thailand)
- 75, 90, 105 and 120 for the Eastern, Central, Rocky Mountain and Pacific time zones (United States)

Note

eq. 31 of reference

References

Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56. FAO, Rome, 300(9).

| | |
|----|--------------------------|
| ws | <i>Sunset hour angle</i> |
|----|--------------------------|

Description

Sunset hour angle for given latitude and solar declination ([SolarDec](#)).

Usage

```
ws(x, lat.rad)
```

Arguments

| | |
|---------|-------------------------------------|
| x | date-time object or day of the year |
| lat.rad | latitude [rad] |

Details

x: must be provided as number (1-366) or as a common date-time object (e.g. POSIXct, POSIXlt or Date objects). All formats for which is.timepoint from the lubridate package returns TRUE can be used

Value

Sunset hour angle

Note

eq. 25 of reference

References

Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). Crop evapotranspiration-Guidelines for computing crop water requirements-FAO Irrigation and drainage paper 56. FAO, Rome, 300(9).

Examples

```
ws(x = 246, lat.rad = -0.35)
```

Index

*Topic **datasets**

constDefaults, 3

controlDefaults, 4

adj_u2, 2, 4

constDefaults, 3, 9–11, 13, 15, 17, 23

controlDefaults, 4, 9–11, 13, 15, 17, 21, 23

deltaVP, 5

dlh, 6

dr, 7

e0, 7, 25, 27

estG, 4, 8, 13

estP, 3, 4, 10, 12, 16

estRs, 4, 11, 12

ETref, 4, 12

prep.date, 15

psyc_cons, 3, 15, 26

Ra, 16

Rn, 4, 9, 18

Rn1, 18, 19, 20

Rns, 18, 19, 22

Rso, 20, 23

satVP, 8, 24, 27

SolarDec, 25, 28

VP, 8, 25, 26

w, 27

ws, 28